

# About periodic solutions of a planar system modelling neural activity\*

S. Treskov, E. Volokitin

## Abstract

We derived explicit symbolic expressions for the first, second, and third Lyapunov coefficients of the complex focus of a planar system modelling activity of a neural network. The analysis of these expressions allowed us to obtain new results about the number and location of limit cycles in the model.

**Introduction.** In this paper we study the mathematical model which describe processes in a neural network consisting from two neurons. The model was suggested in [1] and is defined as a planar differential system

$$\begin{aligned}\dot{u}_1 &= -u_1 + q_{11}\varphi(u_1) - q_{12}u_2 + e_1, \\ \dot{u}_2 &= -u_2 + q_{21}\varphi(u_1) + e_2,\end{aligned}\tag{1}$$

where  $q_{11}, q_{12}, q_{21} > 0$ ,  $e_1, e_2 \in \mathbf{R}$  are parameters, and

$$\varphi(u_1) = \frac{1}{1 + e^{-4u_1}}.$$

In [1] it was fulfilled a study of model (1) with the help of the bifurcation theory and it was listed possible types of dynamical behavior of the neural network; with analytic and numerical techniques it was constructed the bifurcation diagram of system (1) for some parameter values.

In [2] it was proved by analytic methods the correctness of the bifurcation diagram for some specific cases.

---

\*The work is supported by the Russian Foundation for Fundamental Research (Grant No. 99-01-00574)

In [2] it was demonstrated that the change of variables

$$u = u_1, v = \frac{u_2 - e_2}{q_{21}},$$

forms system (1) to the system

$$\begin{aligned}\dot{u} &= -u + a\varphi(u) - bv + c, \\ \dot{v} &= -v + \varphi(u),\end{aligned}\tag{2}$$

which involves three parameters only

$$a = q_{11}, b = q_{12}q_{21}, c = e_1 - q_{12}e_2, a, b > 0, c \in \mathbf{R}.$$

We intend to study system (2).

Using our results from [3], we derived analytic formulas for the first, second, and third Lyapunov coefficients of the complex focus of system (2). A sophisticated treatment of the formulas allows us to assert that for some parameter values system (2) has three concentric limit cycles around a unique steady state, two of these cycles being stable. Such a phase portrait is absent in the list of possible phase portraits of (2) from [1,2] although on our mind it is interesting from a physical point of view.

**1.** Let us to note the following property of system (2). If  $u(t), v(t)$  is the solution of the system then  $u_1(t) = -u(t), v_1(t) = 1 - v(t)$  is the solution of the system

$$\begin{aligned}\dot{u} &= -u + a\varphi(u) - bv - c + b - a, \\ \dot{v} &= -v + \varphi(u),\end{aligned}$$

This fact means that if we have a phase portrait of (2) for parameter values  $a = a_1, b = b_1, c = c_1$  then there exists a symmetric with respect to the point  $u = 0, v = 1/2$  phase portrait of the system for parameter values  $a = a_1, b = b_1, c = c_2 = -a_1 + b_1 - c_1$ . In the parameter space points  $(a_1, b_1, c_1)$  and  $(a_1, b_1, c_2)$  lie on different sides from the hyperplane  $a - b + 2 = 0$  (on equal distances). So, it is sufficiently to study system (2) for parameter values from one of a half-space, for example, from the half-space  $a - b + 2c \geq 0$ .

The mentioned property is obvious from the bifurcation diagrams constructed with numerical methods in [2].

The authors of [1,2] solved all problems of the number, location, and types of steady states of system (2). The system may have between one and three

steady states in the finite part of a phase plane. If the system has three steady states then one of them is always a saddle. Two remaining steady states are nodes or foci (stable or unstable). We distinguish between them a left and right steady state.

The authors of [4] introduced the following method to describe the phase portraits of system (2). The symbol  $s$  ( $u$ ) denotes a stable (unstable) equilibrium; the symbol  $S$  ( $U$ ), a stable (unstable) limit cycle. The subscript 1 refers to the symbol corresponding to the left equilibrium or to a limit cycle around it; the subscript 2, to the symbol corresponding to the right equilibrium or to a limit cycle around it. A symbol has no subscripts when the steady state is unique. The symbols  $S$  and  $U$  without subscripts refer also to limit cycles around all three steady states. If there are several cycles with the same subscript or without subscript, then they are listed in order from inside.

The authors of [1,2] detected following phase portraits of system (2).<sup>1</sup>

|                                    |                                  |                                  |
|------------------------------------|----------------------------------|----------------------------------|
| 1. $s_1 s_2$                       | 5 <sub>1</sub> . $s_1 u_2$       | 9 <sub>1</sub> . $s_1 U_1 u_2 S$ |
| 2. $s_1 s_2 U S$                   | 6. $u_1 s_2 U S$                 | 10. $u_1 s_2 S$                  |
| 3. $s_1 U_1 s_2 U S$               | 6 <sub>1</sub> . $s_1 u_2 U S$   | 10 <sub>1</sub> . $s_1 u_2 S$    |
| 3 <sub>1</sub> . $s_1 s_2 U_2 U S$ | 7. $s_1 U_1 s_2 S$               | 11. $u_1 u_2 S$                  |
| 4. $s_1 U_1 s_2$                   | 7 <sub>1</sub> . $s_1 s_2 U_2 S$ | 12. $s$                          |
| 4 <sub>1</sub> . $s_1 s_2 U_2$     | 8. $s_1 U_1 s_2 U_2 S$           | 13. $s U S$                      |
| 5. $u_1 s_2$                       | 9. $u_1 s_2 U_2 S$               | 14. $u S$                        |

(3)

As a result of our study, we detect one more phase portrait of system (2) which is described by the sequence

$$15. u S U S \tag{4}$$

and corresponds to the situation when in the system two stable limit cycles coexist which are separated with an unstable cycle.

The detection is due to a more comprehensive analysis of the Lyapunov coefficients of the complex focus of system (2). The next section is devoted to this analysis.

---

<sup>1</sup>The subscript 1 for the number of a phase portrait implies that a corresponding portrait is symmetric to the portrait with the same number without the subscript.

2. In [3] we derived symbolic expressions for the first, second, and third Lyapunov coefficients of the complex focus of the system

$$\begin{aligned}\dot{x} &= y, \\ \dot{y} &= a_{10}x + a_{01}y + a_{20}x^2 + a_{11}xy + a_{02}y^2 + \dots\end{aligned}$$

in terms of the coefficients  $a_{ij}$ .

The expressions are rather unwieldy and in general case are not amenable to theoretical study but in some special cases they may be simplified noticeably. In particular, if we have a Liénard system with the complex focus in the origin

$$\begin{aligned}\dot{x} &= y, \\ \dot{y} &= p(x) + yq(x) \equiv p_1x + p_2x^3 + \dots + y(q_1x + q_2x^2 + \dots),\end{aligned}\tag{5}$$

then the expressions for the first three Lyapunov coefficients with accuracy to a positive factor coincide with ones

$$\begin{aligned}l_1 &= p_2q_1 - p_1q_2, \\ l_2 &= 5(p_2q_3 - p_3q_2) + 3(p_4q_1 - p_1q_4), \\ l_3 &= 14p_2(p_2q_4 - p_4q_2) + 21p_1(p_3q_4 - p_4q_3) + 35p_1(p_5q_2 - p_2q_5) + \\ &\quad + 15p_1(p_1q_6 - p_6q_1).\end{aligned}\tag{6}$$

Let  $(u_0, v_0)$  be a steady state of system (2). Then we have

$$u_0 - (a - b)\varphi(u_0) - c = 0, v_0 = \varphi(u_0).$$

Let us make the following change of variables in system (2)

$$x = u - u_0, \quad y = -u + a\varphi(u) - bv + c.$$

After it the steady state  $(u_0, v_0)$  transfers to the origin.

We obtain a Liénard system which is equivalent to system (2)

$$\begin{aligned}\dot{x} &= y, \\ \dot{y} &= p(x) + yq(x),\end{aligned}\tag{7}$$

where

$$\begin{aligned}p(x) &= c + (a - b)/(1 + \exp(-4u_0 - 4x)) - u_0 - x, \\ q(x) &= -2 + (4a \exp(-4u_0 - 4x)/(1 + \exp(-4u_0 - 4x)))^2.\end{aligned}\tag{8}$$

The origin will be the complex focus of system (7), (8) if  $p(0) = q(0) = 0, p'(0) < 0$ , that is under the fulfillment of the conditions

$$\begin{aligned} c + (a - b)/(1 + \exp(-4u_0)) - u_0 &= 0, \\ -2 + (4a \exp(-4u_0)/(1 + \exp(-4u_0))^2 &= 0, \\ -1 + 4(a - b) \exp(4u_0)/(1 + \exp(4u_0))^2 &< 0, \end{aligned}$$

which may be written as

$$\begin{aligned} c &= (b - a)/(1 + \exp(-4u_0)) + u_0, \\ a &= (1 + \exp(4u_0))^2/(2 \exp(4u_0)), \\ -1 + 4(a - b) \exp(4u_0)/(1 + \exp(4u_0))^2 &< 0. \end{aligned} \tag{9}$$

Under conditions (9) system (7), (8) takes form (5).

Evaluating needed coefficients  $p_i, q_i$  and substituting them to (6), we obtain the expressions for the Lyapunov coefficients of system (7), (8).

After some transformations we obtain that the Lyapunov coefficients with accuracy to a positive factor are equal to the expressions

$$\begin{aligned} l_1 &= 1 - 2 \exp(4u_0) - (6 - 8a + 8b) \exp(8u_0) - 2 \exp(12u_0) + \\ &\quad + \exp(16u_0), \\ l_2 &= 3 - 72 \exp(4u_0) + (45 + 8(a - b)) \exp(8u_0) + \\ &\quad (240 - 368(a - b)) \exp(12u_0) + (45 + 8(a - b)) \exp(16u_0) + \\ &\quad - 72 \exp(20u_0) + 3 \exp(24u_0), \\ l_3 &= 1 - 4(29 + a - b) \exp(4u_0) + (717 + 160(a - b)) \exp(8u_0) + \\ &\quad + 16(102 - 205(a - b) - 6(a - b)^2) \exp(12u_0) - \\ &\quad - 2(903 - 1904(a - b) - 384(a - b)^2) \exp(16u_0) + \\ &\quad + 8(651 + 1813(a - b) - 1608(a - b)^2) \exp(20u_0) - \\ &\quad - 2(903 - 1904(a - b) - 364(a - b)^2) \exp(24u_0) + \\ &\quad + 16(102 - 205(a - b) - 6(a - b)^2) \exp(28u_0) + \\ &\quad + (717 + 160(a - b)) \exp(32u_0) - 4(29 + a - b) \exp(36u_0) + \\ &\quad + \exp(40u_0). \end{aligned} \tag{10}$$

Conditions (9) define a two-dimensional bifurcation manifold in the parameter space. This manifold corresponds to the codim 1 Andronov-Hopf bifurcation. The condition  $l_1 = 0$  from (10) defines in this manifold a one-dimensional bifurcation codim 2 manifold corresponding to the degenerate

Andronov-Hopf bifurcation (the Bautin bifurcation, the Takens bifurcation, the vanishing of the first Lyapunov coefficient).

The last manifold is piecewise smooth and it may be defined as follows

$$\begin{aligned} a &= (1 + \vartheta)^2 / (2\vartheta), \\ b &= (1 + \vartheta)^2 (1 + \vartheta^2) / (8\vartheta^2), \\ c &= (1 - 3\vartheta - 3\vartheta^2 + \vartheta^3 + 2\vartheta \ln \vartheta) / (8\vartheta), \\ 0 &< \vartheta < +\infty, \vartheta \neq 1. \end{aligned} \tag{11}$$

Coordinates  $(u_0, v_0)$  of a corresponding unique equilibrium may be obtained from equations  $\exp(4u_0) = \vartheta$ ,  $v_0 = \varphi(u_0)$ .

If parameters  $a, b, c$  are evaluated accordingly to (11) then system (7), (8) (and then system (2) also) has the complex focus of the multiplicity 2. Stability of the focus is defined by the sign of the expression  $\bar{l}_2$  which is the second Lyapunov coefficient  $l_2$  from (10) evaluating accordingly to the (11) and to the equality  $\vartheta = \exp(4u_0)$ .

$$\bar{l}_2 = 2(1 + \vartheta)^2(1 - 14\vartheta + 6\vartheta^2 - 14\vartheta^3 + \vartheta^4).$$

The equation  $\bar{l}_2 = 0$  has two positive roots  $\vartheta_1, \vartheta_2$

$$\begin{aligned} \vartheta_1 &= (7 + 3\sqrt{5} + \sqrt{6}\sqrt{15 + \sqrt{5}}) / 2 \approx 13.6349, \\ \vartheta_2 &= (7 + 3\sqrt{5} - \sqrt{6}\sqrt{15 + \sqrt{5}}) / 2 \approx .0733414, \end{aligned}$$

to which correspond the following parameter values  $a_1 \approx 7.8541$ ,  $b_1 \approx 26.9164$ ,  $c_1 \approx 18.4129$  and  $a_2 = a_1$ ,  $b_2 = b_1$ ,  $c_2 \approx 0.64937$ ,  $a_1 - b_1 + c_1 > 0$ ,  $a_2 - b_2 + c_2 < 0$ .

The third Lyapunov coefficient is negativ for  $(a_1, b_1, c_1)$  and  $(a_2, b_2, c_2)$ .

For such parameters system (2) has a unique steady state which is the complex focus of multiplicity 3.

It is well known that in the parameter space to the point corresponding to the vanishing of the second Lyapunov coefficient is adjacent a region of parameter values for which the system has three concentric limit cycles around the equilibrium.

Thus, we proved that there are parameter values  $a, b, c$ , for which system (2) has three limit cycles surrouding a unique steady state (an unstable focus). Two from these cycles are stable and they are separated by an unstable cycle.

**3.** Fig.1 gives a part of a complete bifurcation diagram of system (2) for a fixed value of parameter  $a > a_1$ . The rest of the diagram may be easily restored from the about mentioned reasons of the symmetry. The diagram is fairly typical. In particular, it has many common features with the diagram given in [1,2] for  $a < a_1$ . We use the notations introduced in [1,2] and restrict ourselves to only brief remarks. More detailed description of the mentioned bifurcations may be found in [6,7], for example.

The curve  $sn_1$  corresponds to the availability in the system of a double steady state which by an appropriate parameter perturbation splits into a saddle and a node which lies left then a saddle. On the curve  $sn_1$  there is a point  $tb_1$ , which corresponds to the Bogdanov-Takens bifurcation and divides the curve into two segments with stable and unstable saddle-nodes.

The line  $h_1$  ( $h_2$ ) corresponds to the Andronov-Hopf bifurcation of the left (right) steady state if parameters are inside the region of multiplicity of steady states or of a unique steady state. As we noted above these lines are defined by formulas (9). The point  $dh_1$  on the line  $h_1$  corresponds to the degenerate Andronov-Hopf bifurcation and divides the line into two segments corresponding to a soft or hard loss in stability. The second Lyapunov coefficient is positive at the point  $dh_1$ . Let us note that this coefficient is negative at the point  $dh_1$  on the diagram presented in [1,2] for  $a < a_1$ .

The curves  $sl_1$  and  $sl_2$  correspond to separatrix loops around left and right steady states. The curve  $sl_4$  corresponds to a separatrix loop around both steady states.

The curve  $snpo$  corresponds to a double cycle which is stable from outside. The curve  $snpo_1$  corresponds to a double cycle which is stable from inside. The curves  $snpo$  and  $snpo_1$  coalesce and form a cusp at the point corresponding to a triple cycle of system (2).

The listed curves divides the parameter space into regions within which the system has distinct rough phase portraits describing in (3), (4).

As usual, we give only the scaleless relative positions of the bifurcation sets. The drawing of these sets to real scale involves technical difficulties.

The local bifurcation sets  $sn_i, h_i, tb_i, dh_i$  can be described by explicit formulas (see (9), (11), for example). Therefore, construction of these sets involves no difficulties. As to sets of nonlocal bifurcations  $sl_i, snpo, snpo_1$ , in constructing them we used the results of the bifurcation theory, and the results of the qualitative theory of ordinary differential equations, and the

results of simulations.

A considerable difference of our bifurcation diagram from the diagram in [1,2] (for  $a < a_1$ ) is the presence of region 15 and curve  $snpo_1$  whose existence we proved above.

In Fig.2 is shown the phase portrait of system (2) for parameter values  $a = 16., b = 130., c = 111.165$ , corresponding to region 15.<sup>2</sup>




Fig.1




Fig.2

Hence our analysis of formulas for the Lyapunov coefficients for the system describing the dynamics of a neural network consisting from two sells enables

---

<sup>2</sup>To confirm the existence of three limit cycles in system (2) for the specified parameter values we studied the proper Poincaré map numerically and established that it has three fixed points.



us to supplement results of other authors. In particular, to earlier described phase portraits we add one more portrait with three limit cycles around an unstable steady state. Studing a complete bifurcation diagram, we specify the position of this portrait in the parameter space.

The results of [1,2] and our results give grounds to advocate that all possible phase portraits of system (2) are included in the join of lists (3), (4). 4. As we noted above, the bifurcation diagram from [1,2] is fairly typical and occurs during a study of many mathematical models, see [8,9] for references. In particular, a similar diagram was consnructed by us in [10] where we investigated a planar system modelling a nerve conduction in the squid giant axon.

**Acknowledgmens.** We thank Professor Fotios Giannakopoulos for materials which he placed at our disposal.

## References

- [1] F. Giannakopoulos, O. Oster. *Bifurcation properties of a planar system modelling neural activity*. Planar Nonlinear Dynamical Systems (Delft, 1995), Differential and Dynamical Systems. 1997. V. 5. 229–242.
- [2] R.E.Kooij, F. Giannakopoulos. *Periodic orbits in planar systems modelling neural activity*. Quarterly of Applied Mathematics. 2000. V.63. No 3. 437–457.
- [3] Volokitin E.P, Treskov S.A. *About the Lyapunov values of the complex focus of a planar dynamical system*. Transactions of Russian Academy of Natural Sciences. 1997. V. 1. No 1. 59–72. (in Russian)
- [4] Vaganov, D.A., Samoilenko, N.G. and Abramov, V.G. *A determination of regions of periodic regimes of continuous stirred tank reactors*. Transactions of Academy of Sciences of the USSR. 1977. V. 234, 640–643. (in Russian)
- [5] Bazykin, A., Kuznetsov, Y. and Khibnik, A. *Portraits of Bifurcations: Bifurcation Diagrams of Planar Dynamical Systems*. 1989. Znanie, Moscow. (in Russian)
- [6] Y.A.Kuznetsov. *Elements of Applied Bifurcation Theory*. 1995. Springer-Verlag, New York.
- [7] A.Khibnik, B.Krauskopf, C.Rousseau. *Global study of a family of cubic Lienard equations*. Preprint. CRM-2471, April 1997.
- [8] A.Khibnik, B.Krauskopf, C. Rousseau. *Global study of a family of cubic Lienard equations*. Nonlinearity. 1998. V.11. No 6, 1505–1519.
- [9] Volokitin E.P, Treskov S.A. *A parametric portrait of the Fitz-Hugh system*. Mathematical modelling. 1994. No 6. 65–78. (in Russian)